



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Therefore, the only possible solutions are

$$xu - 5yv = 2, \quad xu + 5yv = \pm 2.$$

Whence, $2xu = 4$ or 0 , $10yv = 0$ or -4 ; i. e., either x , u , y , or v equals zero.

Solved, similarly, by G. B. M. Zerr and V. M. Spunar.

II. Solution by O. C. CARMICHAEL, Oxford, Ala.

If the square of the first equation be added to five times the square of the second equation, we have

$$x^2u^2 + 5x^2v^2 + 25y^2v^2 + 5y^2u^2 = 9.$$

Therefore, there is no integral solution in x , y , u , v except when one of the unknowns is zero; for, if they were all positive integers, $25y^2v^2$ itself would be greater than 9.

303. Proposed by PROF. R. D. CARMICHAEL, Anniston, Ala.

Evaluate the determinant

$$\begin{vmatrix} D_1 & x_1x_2 & x_1x_3 & \dots & x_1x_n \\ x_1x_2 & D_2 & x_2x_3 & \dots & x_2x_n \\ x_1x_3 & x_2x_3 & D_3 & \dots & x_3x_n \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ x_1x_n & x_2x_n & x_3x_n & \dots & D_n \end{vmatrix}$$

Solution by J. W. CLAWSON, Ursinus College, Collegeville, Pa.

$$\begin{aligned} \Delta &= x_1x_2x_3\dots x_n \begin{vmatrix} D_1/x_1 & x_2 & x_3 & \dots & x_n \\ x_1 & D_2/x_2 & x_3 & \dots & x_n \\ x_1 & x_2 & D_3/x_3 & \dots & x_n \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ x_1 & x_2 & x_3 & \dots & D_n/x_n \end{vmatrix} \\ &= \prod_{i=1}^n x_i \begin{vmatrix} (D_1 - x_1^2)/x_1 & 0 & 0 & \dots & (x_n^2 - D_n)/x_n \\ 0 & (D_2 - x_2^2)/x_2 & 0 & \dots & (x_n^2 - D_n)/x_n \\ 0 & 0 & (D_3 - x_3^2)/x_3 & \dots & (x_n^2 - D_n)/x_n \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ x_1 & x_2 & x_3 & \dots & D_n/x_n \end{vmatrix} \end{aligned}$$

Subtracting the last row from each row,

$$= \prod_{r=1}^n x_r \cdot \prod_{r=1}^n \frac{D_r - x_r^2}{x_r} \begin{vmatrix} 1 & 0 & 0 & \cdots & -1 \\ 0 & 1 & 0 & \cdots & -1 \\ 0 & 0 & 1 & \cdots & -1 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \frac{x_1^2}{D_1 - x_1^2} & \frac{x_2^2}{D_2 - x_2^2} & \frac{x_3^2}{D_3 - x_3^2} & \cdots & \frac{D_n}{D_n - x_n^2} \end{vmatrix}$$

$$= \prod_{r=1}^n (D_r - x_r^2) \begin{vmatrix} 1 & 0 & 0 & \cdots & 0 \\ 0 & 1 & 0 & \cdots & -1 \\ 0 & 0 & 1 & \cdots & -1 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \frac{x_1^2}{D_1 - x_1^2} & \frac{x_2^2}{D_2 - x_2^2} & \frac{x_3^2}{D_3 - x_3^2} & \cdots & \frac{D_n}{D_n - x_n^2} + \frac{x_1^2}{D_1 - x_1^2} \end{vmatrix}$$

adding last column to the first. Since the first row contains $n-1$ zeros, this reduces the determinant to one of the $(n-1)$ st order. Adding last column to the first we reduce to one of the $(n-2)$ nd order. Repeat this process $(n-1)$ times in all.

$$\begin{aligned} \text{Then } \Delta &= \prod_{r=1}^n (D_r - x_r^2) \left[\frac{D_n}{D_n - x_n^2} + \frac{x_1^2}{D_1 - x_1^2} + \frac{x_2^2}{D_2 - x_2^2} + \cdots + \frac{x_{n-1}^2}{D_{n-1} - x_{n-1}^2} \right] \\ &= \prod_{r=1}^n (D_r - x_r^2) \left[\frac{x_1^2}{D_1 - x_1^2} + \frac{x_2^2}{D_2 - x_2^2} + \cdots + \frac{x_{n-1}^2}{D_{n-1} - x_{n-1}^2} + \frac{x_n^2}{D_n - x_n^2} + 1 \right] \\ &= \prod_{r=1}^n (D_r - x_r^2) \left[\sum_{r=1}^n \frac{x_r^2}{D_r - x_r^2} + 1 \right]. \end{aligned}$$

Also solved by G. B. M. Zerr, V. M. Spunar, and J. Scheffer.

304. Proposed by C. N. SCHMALL, New York City.

A policeman on a motor-cycle starts in pursuit of an automobile when the latter has a headway of $\frac{1}{2}$ a mile. A pedestrian who is $\frac{1}{4}$ mile ahead of the auto and who is walking at the rate of 5 miles an hour, notices that when the auto overtakes him the policeman is only $5-12$ of a mile behind the auto, and $2\frac{1}{2}$ miles from where the officer started; he overtakes the auto. How long did the chase last?

Solution by G. B. M. ZERR, A. M., Ph. D., and the PROPOSER.

Let x =policeman's rate, y =auto's rate, and z =time for auto to overtake pedestrian.

$$\text{Then } (5z + \frac{1}{4})/y = (5z + \frac{1}{4} + \frac{1}{12})/x \dots (1),$$

$$2/y = (2\frac{1}{2})/x \dots (2).$$

$$(1)/(2) \text{ gives } 300z + 15 = 240z + 16. \quad \therefore z = \frac{1}{60} \text{ hours} = 1 \text{ minute.}$$